

Complex Analysis as Catalyst

Steven G. Krantz¹

Abstract: We see the subject of complex analysis, in both one and several variables, as an excuse to study other areas: real variables, topology, group theory, abstract algebra, partial differential equations, or geometry. The purpose of this paper is to develop that theme, illustrated by several examples.

0 Prefatory Thoughts

I am known as a complex analyst, yet I have never had a course in complex variable theory. This lacuna in my education has perhaps had an effect on the way that I view the subject.

I see the elegant subject of complex function theory as a stage for the setting of beautiful problems. Once these problems are apprehended or understood, they are often best solved by stripping away the complex variables and bringing in tools from other parts of mathematics. That is the point of view that we promulgate in the present article. We illustrate the point with examples from real analysis, group theory, abstract algebra, partial differential equations, geometry, and other subjects as well.

1 An Example from Topology

Topology has become part of the bedrock of modern mathematics. Originally studied as *analysis situs*, it now permeates all branches of our discipline. We understand that the entire texture of a subject may be influenced by the particular topology that is chosen. The examples in this section illustrate the point.

In this paper a *domain* will be a connected, open set in $\mathbb{C}$ or $\mathbb{C}^n$. Let Ω be a domain. It is a commonplace in complex variable theory to introduce the

¹We are happy to thank the American Institute of Mathematics for its hospitality and support during this work.

notion of *uniform convergence on compact subsets* of Ω . Though not often noted, this is in fact equivalent to the compact-open topology on the space $\mathcal{O}(\Omega)$ of holomorphic functions on Ω . In practice we say that $f_j \in \mathcal{O}(\Omega)$ converge uniformly on compact subsets of Ω to a function $f \in \mathcal{O}(\Omega)$ if, for any compact $K \subseteq \Omega$ and any $\epsilon > 0$, there is a $J > 0$ such that if $j > J$ then $|f_j(z) - f(z)| < \epsilon$ for all $z \in K$. The basic result, which every student learns, is that the limit function f is holomorphic on Ω provided only that the elements f_j are holomorphic on Ω .

An interesting and natural question to ask at this point is

What happens if the $f_j \in \mathcal{O}(\Omega)$ only converge pointwise?

A naive answer might be “Nothing much,” but that would be hasty and also incorrect. Certainly the limit function f is well defined and we may ask what properties it has. Is it continuous? Probably not. It turns out that the correct statement is this:

Theorem 1 *Let $f_j \in \mathcal{O}(\Omega)$ converge pointwise to a limit function f on Ω . Then there is a dense, open subset $V \subseteq \Omega$ such that f is holomorphic on V .*

As one might suppose from the statement, the proof will involve the Baire category theorem. We see then, in our first example, that even though the theorem is about function theory the proof will consist of real analysis and topology.

Proof of Theorem 1: Let U be an open subset of Ω with compact closure in Ω . Define, for $k = 1, 2, \dots$,

$$S_k = \{z \in \overline{U} : |f_j(z)| \leq k \text{ for all } j \in \mathbb{N}\}.$$

Since the f_j converge at each $z \in \overline{U}$, certainly the set $\{f_j(z)\}$ is bounded for each fixed z . So each $z \in \overline{U}$ lies in some S_k . In other words,

$$\overline{U} = \bigcup_k S_k.$$

Now of course $\overline{U}$ is a complete metric space (in the ordinary Euclidean metric), so the Baire category theorem tells us that some S_k must be “somewhere dense” in $\overline{U}$. This means that $\overline{S_k}$ will contain a nontrivial Euclidean metric ball in $\overline{U}$. Call the ball B . Now it is a simple matter to apply Montel’s

theorem on B to find a subsequence f_{j_k} that converges uniformly on compact sets to a limit function g . But of course g must coincide with f , and g (hence f) must be holomorphic on B .

Since the choice of U in the above arguments was arbitrary, the conclusion of the theorem follows. $\square$

One might ask what more can be said about the open, dense set on which the limit function f is holomorphic. Put in other words, what can one say about $\Omega \setminus V$? In fact Siciak [SIC] has given a rather thorough answer to this question. The statement and proofs involve analytic capacity, and we cannot treat the details here.

2 Hard Analysis

The central feature of the basic theory of complex analysis is the Cauchy integral formula. A basic form of this result is as follows:

Proposition 2 *Let f be holomorphic in a neighborhood of the closed disc $\overline{D}(P, r)$ in the complex plane. If $z \in D(P, r)$ then*

$$f(z) = \frac{1}{2\pi i} \oint_{\partial D(P, r)} \frac{f(\zeta)}{\zeta - z} d\zeta. \quad (*)$$

A reasonable—though not often formulated—question to ask is, “What is so special about holomorphic functions? Is the formula (*) true only for holomorphic functions? Is something analogous true for a more general class of functions? Is there a universal formula of which (*) is just a special case?”

The somewhat surprising answer to these questions is “yes”. We shall, in the present section, prove that result. Later sections will give applications of the more general integral formula.

The experienced reader will know that the Poisson integral formula is an identity that is closely related to (*) (see [GRK], [KRA4], [KRA5]), and which is valid for harmonic functions. So it can properly be considered a generalization of the Cauchy formula. But that is not what we seek at this time. First of all, it is too well known hence not suitable grist for our mill. Secondly, we seek a formula that is valid for all smooth functions. In fact our result is this:

Theorem 3 . Let $\Omega \subseteq \mathbb{C}$ be a bounded domain with C^1 boundary (i.e., the boundary consists of finitely many closed, C^1 curves). Let f be a continuously differentiable function on $\overline{\Omega}$. Then, for any $z \in \Omega$,

$$f(z) = \frac{1}{2\pi i} \oint_{\partial\Omega} \frac{f(\zeta)}{\zeta - z} d\zeta - \frac{1}{2\pi i} \iint_{\Omega} \left(\frac{\partial f / \partial \bar{\zeta}}{\zeta - z} \right) d\bar{\zeta} \wedge d\zeta .$$

[In the proof we shall review the notation $\partial\partial\bar{\zeta}$.]

Our proof will use a form of Stokes's theorem that may be unfamiliar. Recall that the classical Stokes's theorem that can be found in any calculus book (see, for example, [BLK]) says that, if u_1, u_2 are continuously differentiable functions on $\overline{\Omega}$, then

$$\int_{\partial\Omega} u_1 dx + \int_{\partial\Omega} u_2 dy = \iint_{\Omega} \frac{\partial u_2}{\partial x} - \frac{\partial u_1}{\partial y} dx \wedge dy . \quad (\star)$$

Note that, in complex notation,

$$\begin{aligned} \frac{\partial}{\partial z} &= \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) , \\ \frac{\partial}{\partial \bar{z}} &= \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) , \end{aligned}$$

$dz = dx + idy$, and $d\bar{z} = dx - idy$. Thus a simple change of variable in the formula $(\star)$ yields the “complex” version of Stokes's theorem

$$\oint_{\partial\Omega} u dz = \iint_{\Omega} \frac{\partial u}{\partial \bar{z}} d\bar{z} \wedge dz . \quad (\star\star)$$

We shall make good use of this version of Stokes's result in the argument that follows.

Proof of Theorem 3: We shall unashamedly use Stokes's theorem, and a mild dose of the theory of differential forms. This treatment will be self-contained, and even the nervous reader should find it enlightening.

Fix a point $z \in \Omega$ and a positive number ϵ that is less than the Euclidean distance of z to $\partial\Omega$. Define

$$\Omega_\epsilon \equiv \Omega \setminus \overline{D}(z, \epsilon) .$$

We apply the aforementioned version of Stokes's theorem to the function

$$g(\zeta) \equiv \frac{f(\zeta)}{\zeta - z}$$

on the domain Ω_ϵ .

The result is

$$\oint_{\partial\Omega_\epsilon} g(\zeta) d\zeta = \iint_{\Omega_\epsilon} \frac{\partial}{\partial\bar{\zeta}} (g(\zeta)) d\bar{\zeta} \wedge d\zeta.$$

Writing out the definition of g gives

$$\oint_{\partial\Omega_\epsilon} \frac{f(\zeta)}{\zeta - z} d\zeta = \iint_{\Omega_\epsilon} \frac{\partial}{\partial\bar{\zeta}} \left(\frac{f(\zeta)}{\zeta - z} \right) d\bar{\zeta} \wedge d\zeta$$

or

$$\oint_{\partial\Omega_\epsilon} \frac{f(\zeta)}{\zeta - z} d\zeta = \iint_{\Omega_\epsilon} \left(\frac{\partial f / \partial\bar{\zeta}}{\zeta - z} \right) d\bar{\zeta} \wedge d\zeta. \quad (\dagger)$$

We may write out the lefthand side of $(\dagger)$ as

$$\oint_{\partial\Omega} \frac{f(\zeta)}{\zeta - z} d\zeta - \oint_{\partial D(z, \epsilon)} \frac{f(\zeta)}{\zeta - z} d\zeta \equiv I + II.$$

Note the differing orientations of the two pieces of the boundary. We examine II :

$$\begin{aligned} II &= \int_0^{2\pi} \frac{f(z + \epsilon e^{it})}{\epsilon e^{it}} \cdot i\epsilon e^{it} dt \\ &= i \int_0^{2\pi} f(z + \epsilon e^{it}) dt. \end{aligned}$$

A simple limiting argument, using the continuity of f , shows that this last line tends as $\epsilon \rightarrow 0^+$ to $2\pi i f(z)$.

Thus we have from $(\dagger)$ (letting $\epsilon \rightarrow 0^+$ on the right as well) that

$$\oint_{\partial\Omega} \frac{f(\zeta)}{\zeta - z} d\zeta - 2\pi i f(z) = \iint_{\Omega} \left(\frac{\partial f / \partial\bar{\zeta}}{\zeta - z} \right) d\bar{\zeta} \wedge d\zeta.$$

In conclusion,

$$f(z) = \frac{1}{2\pi i} \oint_{\partial\Omega} \frac{f(\zeta)}{\zeta - z} d\zeta - \frac{1}{2\pi i} \iint_{\Omega} \left(\frac{\partial f / \partial\bar{\zeta}}{\zeta - z} \right) d\bar{\zeta} \wedge d\zeta.$$

That is the desired conclusion. $\square$

Of course it is incumbent on us to point out that, in the case that f is holomorphic, it follows immediately (from the Cauchy-Riemann equations) that $\partial f / \partial \bar{\zeta} \equiv 0$. Hence the formula in the statement of the theorem becomes

$$f(z) = \frac{1}{2\pi i} \oint_{\partial\Omega} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

This is of course the classical Cauchy integral formula.

3 Partial Differential Equations

One of the most profound facts in basic function theory is the Riemann mapping theorem:

Theorem 4 *Let Ω be any simply connected domain in $\mathbb{C}$ that is not the entire complex plane. Then there is a one-to-one, onto conformal mapping of Ω to the unit disc D .*

This result has exerted a profound influence on our subject ever since its original formulation (and not-quite-right) proof. It has been generalized to Koebe’s uniformization theorem, and has inspired the definition of the Carathéodory and Kobayashi metrics (see, for instance, [KRA1] and also our Section 7). If $\varphi : \Omega \rightarrow D$ is the Riemann mapping, then it is natural to think that the function theory on Ω may be transferred to function theory on D by way of φ . This process has many advantages, for D has great symmetry, it is possessed of a transitive automorphism group, and its boundary carries the full arsenal of Fourier analysis.

But many of the most basic questions that one might wish to ask would necessitate that φ be well behaved up to the boundary of Ω . Thus one might ask whether φ and φ^{-1} extend continuously to $\partial\Omega$ and ∂D respectively. Carathéodory has shown (see the details in [GRK]) that, in case $\partial\Omega$ is a Jordan curve, the answer is “yes”. More generally, one might ask whether, if $\partial\Omega$ is C^k (i.e., consists of the union of finitely many C^k curves), then do φ and φ^{-1} extend C^k to the respective boundaries? Phrased slightly differently, we may ask whether φ and φ^{-1} extend to C^k diffeomorphisms of the closures of Ω and D .

These last are profound questions, first studied in the nineteenth century by Painlevé in his Paris thesis. Painlevé's original techniques were purely function-theoretic, but the modern approach to the matter is by way of partial differential equations. A full treatment of the matter requires extensive exposition (see the monograph [KRA2]), but we may indicate the main points here.

In fact the approach that we wish to present requires a detour into the study of the celebrated Dirichlet problem. Now let $\Omega \subseteq \mathbb{C}$ be a bounded domain with C^1 boundary. Let f be a given continuous function on $\partial\Omega$. The *Dirichlet problem* is the elliptic boundary value problem given by

$$\begin{cases} \Delta u &= 0 & \text{on } \Omega \\ u &= f & \text{on } \partial\Omega. \end{cases}$$

As a classical instance, if Ω is a metal sheet of heat-conducting material, and if f represents an initial heat distribution on the boundary of Ω , then the solution u of the Dirichlet problem is the steady state heat distribution over Ω .

There is an extensive theory—due to O. Perron (see [GRK])—for constructing the solution of the Dirichlet problem. In the case that Ω is the unit disc, the Poisson integral formula gives an explicit solution to the Dirichlet problem:

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\psi}) \frac{1-r^2}{1-2r\cos(\theta-\psi)+r^2} d\psi.$$

One notes immediately in this formula that the kernel, that is to say the expression

$$P(r, \theta - \psi) \equiv \frac{1-r^2}{1-2r\cos(\theta-\psi)+r^2},$$

is real analytic in its arguments. It follows then that any harmonic function is real analytic, so certainly smooth. The key question is

What is the behavior of the solution u of the Dirichlet problem
up to the boundary of Ω ?

Perron's original theory shows that, under mild regularity conditions on $\partial\Omega$, the solution to the Dirichlet problem is continuous on the closure. That is to say, the function

$$U(z) = \begin{cases} u(z) & \text{if } z \in \Omega \\ f(z) & \text{if } z \in \partial\Omega \end{cases}$$

is continuous on $\overline{\Omega}$. This assertion may be proved by direct estimation from the Poisson integral formula in the case that Ω is the disc (see [KRA3, Chapter 8]). The higher (or C^k) regularity theory—commonly known as the *Schauder estimates*—is deep and difficult. It is challenging even to find a source that expositis the matter completely and clearly.² We can only content ourselves here with a brief statement (no proof) of the regularity theory for the Laplacian.

First some terminology. Recall that if g is a function on a domain $E \subseteq \mathbb{R}^N$ then we say that g satisfies a *classical Lipschitz condition* if there is a constant $C > 0$ such that

$$|g(x) - g(y)| \leq C \cdot |x - y|$$

for all $x, y \in E$. We sometimes call this a Lipschitz condition of *order 1*. More generally, if $0 < \alpha < 1$, then we say that g satisfies a *Lipschitz condition of order α* provided that

$$|g(x) - g(y)| \leq C \cdot |x - y|^\alpha$$

for all $x, y \in E$.³ Clearly a Lipschitz condition of order 1 is stronger than a Lipschitz condition of order α , $0 < \alpha < 1$.

Now let g be a function on an open domain $U \subseteq \mathbb{R}^N$. Let k be a non-negative integer and $0 < \alpha < 1$. We say that $g \in C^{k,\alpha}$ provided that g is k -times continuously differentiable on U and further that any k^{th} derivative $D^\beta g$ (with β a multi-index of order k) is Lipschitz⁴ of order α . The $C^{k,\alpha}$ form a scale of spaces that stratify C^∞ . Clearly, for example, for fixed α ,

$$\bigcap_k C^{k,\alpha} = C^\infty.$$

Now we may state our main regularity result for the Dirichlet problem for the Laplacian. Again, references are [KRA2], [GIT], and [HOR].

Proposition 5 *Let k be a nonnegative integer. Let $\Omega \subseteq \mathbb{C}$ be a bounded domain with C^k boundary. Let f be a C^k function on $\partial\Omega$. Then the solution u*

²The book [KRA2] is one such reference; see also the classical references [HOR] and [GIT].

³It seems to be the case that harmonic analysts like to say “Lipschitz space” while partial differential equations theorists like to say “Hölder space”. We adhere to the former paradigm.

⁴This definition is easily extended to the situation where the domain of the function is the closure of an open domain.

on Ω of the Dirichlet problem with boundary data f satisfies $u \in C^{(k-1),\alpha}(\bar{\Omega})$ for any $0 < \alpha < 1$.

The fact that we may not assert in this last proposition that $u \in C^k(\bar{\Omega})$ is an unfortunate artifact of the theory of singular integrals (again see [KRA2] and also [STE]). A sharper result may be formulated if we use Sobolev spaces or Besov spaces or Triebel-Lizorkin spaces. One may even formulate a more refined definition of Lipschitz spaces that results in a sharper statement. But we cannot indulge in these technical niceties here.

We next need the following sharp form of the classical lemma of Hopf (which was originally formulated to study the maximum principle for solutions of elliptic partial differential equations):

Lemma 6 (Hopf) *Let $\Omega \subset \mathbb{R}^N$ have C^2 boundary. Let $u \in C(\bar{\Omega})$ with u harmonic and non-constant on Ω . Let $P \in \bar{\Omega}$ and assume that u takes a local minimum at P . Then*

$$\frac{\partial u}{\partial \nu}(P) < 0.$$

Here $\partial/\partial \nu$ is the unit outward normal derivative.

Remark: That the indicated normal derivative is nonpositive is obvious just from examining the Newton quotients. The interesting thing is that the derivative must be strictly negative. $\square$

Proof: Suppose without loss of generality that $u > 0$ on Ω near P and that $u(P) = 0$. Let B_R be a ball that is internally tangent to $\bar{\Omega}$ at P . We may assume that the center of this ball is at the origin and that P has coordinates $(R, 0, \dots, 0)$. Then, by Harnack's inequality (see [GRK]), we have for $0 < r < R$ that

$$u(r, 0, \dots, 0) \geq c \cdot \frac{R^2 - r^2}{R^2 + r^2}$$

hence

$$\frac{u(r, 0, \dots, 0) - u(R, 0, \dots, 0)}{r - R} \leq -c' < 0.$$

Therefore

$$\frac{\partial u}{\partial \nu}(P) \leq -c' < 0.$$

This is the desired result. $\square$

Proposition 7 *Let $\Omega \subseteq \mathbb{C}$ be a bounded, simply connected domain with C^k boundary. Let $\varphi : \Omega \rightarrow D$ be the conformal mapping provided by the Riemann mapping theorem. Then φ extends to a C^{k-2} , univalent, mapping of $\overline{\Omega}$ to $\overline{D}$ for any $0 < \alpha < 1$.*

Proof: Let W be a collared neighborhood of $\partial\Omega$ with smooth boundary. Set $\partial\Omega' = \partial W \cap \Omega$ and let $\partial D' = \varphi(\partial\Omega')$. Define B to be the region bounded by ∂D and $\partial D'$. Let U be the region bounded by $\partial\Omega$ and $\partial\Omega'$. We solve the Dirichlet problem on B with boundary data

$$f(\zeta) = \begin{cases} 1 & \text{if } \zeta \in \partial D \\ 0 & \text{if } \zeta \in \partial D' \end{cases}$$

Call the solution u .

Consider $v \equiv u \circ \varphi : \Omega \rightarrow \mathbb{R}$. Then of course v is still harmonic. By Carathéodory's theorem, v extends to $\partial\Omega, \partial\Omega'$, and

$$v = \begin{cases} 1 & \text{if } \zeta \in \partial\Omega \\ 0 & \text{if } \zeta \in \partial\Omega' \end{cases}$$

The function v will be $C^{(k-1),\alpha}$, $0 < \alpha < 1$, on $\overline{U}$ by Proposition 5. If we consider a first order derivative $\mathcal{D}$ of v we obtain

$$|\mathcal{D}v| = |\mathcal{D}(u \circ \varphi)| = |\nabla u| |\nabla \varphi| \leq C.$$

It follows that

$$|\nabla \varphi| \leq \frac{C}{|\nabla u|}. \quad (*)$$

Now let us return to the u from the Dirichlet problem that we considered prior to line (*). Hopf's lemma tells us that $|\nabla u| \geq c' > 0$ near ∂D . Thus, from (*), we conclude that

$$|\nabla \varphi| \leq C. \quad (**)$$

Thus we have bounds on the first derivatives of φ .

To control the second derivatives, we calculate that

$$\begin{aligned} C &\geq |\nabla^2 v| = |\nabla(\nabla v)| = |\nabla(\nabla(u \circ \varphi))| \\ &= |\nabla(\nabla u(\varphi) \cdot \nabla \varphi)| = |(\nabla^2 u \cdot [\nabla \varphi]^2) + (\nabla u \cdot \nabla^2 \varphi)|. \end{aligned}$$

Here the reader should think of ∇ as representing a generic first derivative and ∇^2 a generic second derivative. We conclude that

$$|\nabla u| |\nabla^2 \varphi| \leq C + |\nabla^2 u| |(\nabla \varphi)^2| \leq C'.$$

Hence (again using Hopf's lemma)

$$|\nabla^2 \varphi| \leq \frac{C}{|\nabla u|} \leq C''.$$

In the same fashion, we may prove that $|\nabla^j \varphi| \leq C_j$, any $0 \leq j \leq k-1$. This means (use the fundamental theorem of calculus) that $\varphi \in C^{k-2}(\bar{\Omega})$. $\square$

We note that several small technical emendations in the proof presented would suffice to show that $\varphi \in C^{(k-1),\alpha}(\bar{\Omega})$ for any $0 < \alpha < 1$. We omit those details.

4 Algebra

Now we turn to a result of Lipman Bers that has a different flavor. It characterizes domains in terms of a certain algebraic invariant.

Let $\Omega \subseteq \mathbb{C}$ be a domain. Let $\mathcal{O}(\Omega)$ denote the algebra of holomorphic functions from Ω to $\mathbb{C}$. Bers's theorem says, in effect, that the algebraic structure of $\mathcal{O}(\Omega)$ characterizes Ω . We begin our study by introducing a little terminology.

Definition 1 Let $\Omega \subseteq \mathbb{C}$ be a domain. A $\mathbb{C}$ -algebra homomorphism $\varphi : \mathcal{O}(\Omega) \rightarrow \mathbb{C}$ is called a *character* of $\mathcal{O}(\Omega)$. If $c \in \mathbb{C}$, then the mapping

$$\begin{aligned} e_c : \mathcal{O}(\Omega) &\rightarrow \mathbb{C}, \\ f &\mapsto f(c), \end{aligned}$$

is called a *point evaluation*. Every point evaluation is a character.

It should be noted that if $\Omega, \hat{\Omega}$ are domains and $\varphi : \mathcal{O}(\Omega) \rightarrow \mathcal{O}(\hat{\Omega})$ is not the trivial zero homomorphism, then $\varphi(1) = 1$. This follows because $\varphi(1) = \varphi(1 \cdot 1) = \varphi(1) \cdot \varphi(1)$. On any open set where the holomorphic

function $\varphi(1)$ does not vanish, we find that $\varphi(1) \equiv 1$. The result follows by analytic continuation.

It turns out that every character of $\mathcal{O}(\Omega)$ is a point evaluation. That is the content of the next lemma.

Lemma 8 *Let φ be a character on $\mathcal{O}(\Omega)$. Then $\varphi = e_c$ for some $c \in \Omega$. Indeed, $c = \varphi(\text{id}) \in \Omega$. Here id is defined by $\text{id}(z) = z$.*

Proof: Let c be defined as in the statement of the lemma. Let $f(z) = z - c$. Then

$$\varphi(f) = \varphi(\text{id}) - \varphi(c) = c - c = 0.$$

If it were not the case that $c \in \Omega$ then the function f would be a unit in $\mathcal{O}(\Omega)$. But then

$$1 = \varphi(f \cdot f^{-1}) = \varphi(f) \cdot \varphi(f^{-1}) = 0.$$

That is a contradiction. So $c \in \Omega$.

Now let $g \in \mathcal{O}(\Omega)$ be arbitrary. Then we may write

$$g(z) = g(c) + f(z) \cdot \tilde{g}(z),$$

where $\tilde{g} \in \mathcal{O}(\Omega)$. Thus

$$\varphi(g) = \varphi(g(c)) + \varphi(f) \cdot \varphi(\tilde{g}) = g(c) + 0 = g(c) = e_c(g).$$

We conclude that $\varphi = e_c$, as was claimed. □

Now we may state and prove Bers's theorem.

Theorem 9 *Let $\Omega, \tilde{\Omega}$ be domains. Suppose that*

$$\varphi : \mathcal{O}(\Omega) \rightarrow \mathcal{O}(\tilde{\Omega})$$

is a $\mathbb{C}$ -algebra homomorphism. Then there exists one and only one holomorphic mapping $h : \tilde{\Omega} \rightarrow \Omega$ such that

$$\varphi(f) = f \circ h \quad \text{for all } f \in \mathcal{O}(\Omega).$$

In fact, the mapping h is given by $h = \varphi(\text{id})$.

The homomorphism φ is bijective if and only if h is conformal, that is, a one-to-one and onto holomorphic mapping from $\tilde{\Omega}$ to Ω .

Proof: Since we want the mapping h to satisfy $\varphi(f) = f \circ h$ for all $f \in \mathcal{O}(\Omega)$, it must in particular satisfy $\varphi(\text{id}_\Omega) = \text{id}_\Omega \circ h = h$. We take this as our definition of the mapping h .

If $a \in \tilde{\Omega}$, then $e_a \circ \varphi$ is a character of $\mathcal{O}(\Omega)$. Thus our lemma tells us that $e_a \circ \varphi$ must in fact be a point evaluation on Ω . As a result,

$$e_a \circ \varphi = e_c, \quad \text{with } c = (e_a \circ \varphi)(\text{id}_\Omega) = e_a(h) = h(a).$$

Thus, if $f \in \mathcal{O}(\Omega)$, then

$$\varphi(f)(a) = e_a(\varphi \circ f) = (e_a \circ \varphi)(f) = e_{h(a)}(f) = f(h(a)) = (f \circ h)(a)$$

for all $a \in \tilde{\Omega}$. We conclude that $\varphi(f) = f \circ h$ for all $f \in \mathcal{O}(\Omega)$.

For the last statement of the theorem, suppose that h is a one-to-one, onto conformal mapping of $\tilde{\Omega}$ to Ω . If $g \in \mathcal{O}(\Omega)$, then set $f = g \circ h^{-1}$. It follows that $\varphi(f) = f \circ h = g$. Hence φ is onto. Likewise, if $\varphi(f_1) = \varphi(f_2)$, then $f_1 \circ h = f_2 \circ h$ hence, composing with h^{-1} , $f_1 \equiv f_2$. So φ is one-to-one. Conversely, suppose that φ is an isomorphism. Let $a \in \Omega$ be arbitrary. Then e_a is a character on $\mathcal{O}(\Omega)$; hence $e_a \circ \varphi^{-1}$ is a character on $\mathcal{O}(\tilde{\Omega})$. By the lemma, there is a point $c \in \tilde{\Omega}$ such that $e_a \circ \varphi^{-1} = e_c$. It follows that

$$e_a = e_c \circ \varphi.$$

Applying both sides of this last identity to id_Ω yields

$$e_a(\text{id}_\Omega) = (e_c \circ \varphi)(\text{id}_\Omega).$$

Unraveling the definitions gives

$$a \equiv e_c(\text{id}_\Omega \circ h) = h(c).$$

Thus $h(c) = a$ and h is surjective. The argument in fact shows that the pre-image c is uniquely determined. So h is also one-to-one. $\square$

As an application⁵ of Bers's theorem, we can see immediately that the disk $D = \{z \in \mathbb{C} : |z| < 1\}$ and the annulus $\mathcal{A} = \{z \in \mathbb{C} : 1/2 < |z| < 2\}$

⁵Of course this application is in a certain sense trivial because the two domains have different topology. All we are doing here is illustrating the algebraic ideas with a different proof.

are not conformally equivalent. For the algebra $\mathcal{O}(D)$ can be generated by 1 and z . But the algebra $\mathcal{O}(\mathcal{A})$ cannot be generated by 1 and just one other function (because natural generators for $\mathcal{O}(\mathcal{A})$ are $\{1, z, 1/z\}$ and it is impossible to come up with a shorter list). We leave the details of these assertions to the reader.

It must be noted, of course, that if Ω_1 and Ω_2 are conformally equivalent via a mapping $\varphi : \Omega_1 \rightarrow \Omega_2$, then their respective algebras of holomorphic functions are trivially isomorphic by way of the conjugation

$$\mathcal{O}(\Omega_1) \ni \eta \longmapsto \varphi \circ \eta \circ \varphi^{-1} \in \mathcal{O}(\Omega_2).$$

The statement of the theorem is the converse of this trivial result, and is definitely more interesting. As an illustration, let $\Omega_1 = D$, the unit disc, and $\Omega_2 = \mathbb{C}$, the complex plane. Of course these domains cannot be conformally equivalent—by a simple application of Liouville’s theorem. We may conclude from Theorem 9 that $\mathcal{O}(D)$ and $\mathcal{O}(\mathbb{C})$ are inequivalent as algebras. This last statement is rather interesting, and is certainly nonobvious. Both algebras can be thought of as the set of all power series

$$\sum_j a_j z^j,$$

and the only difference between the two is the radius of convergence. It is not so clear why the two algebras should be nonisomorphic.

5 Group Theory

We now pass to the subject of the function theory of several complex variables. One of the great classical results was motivated by the Riemann mapping theorem from one complex variable. That venerable result tells us that, in a certain sense, the disc is the “canonical domain” in $\mathbb{C}$. Any simply connected domain is conformally equivalent to the unit disc. Thus one may ask what is the analogous result in several complex variables. What is the canonical domain in $\mathbb{C}^n$?

Two natural candidates for this canonical domain are the unit ball

$$B = \{z = (z_1, z_2, \dots, z_n) \in \mathbb{C}^n : \sum_j |z_j|^2 < 1\}$$

and the unit polydisc

$$D^n = \{z = (z_1, z_2, \dots, z_n) \in \mathbb{C}^n : |z_j| < 1 \text{ for } j = 1, 2, \dots, n\}.$$

Poincaré caused quite a stir in 1906 when he proved that, when $n > 1$, B and D^n are *not* biholomorphically equivalent. That is to say, there does *not* exist a holomorphic mapping

$$\Phi : B \rightarrow D^n$$

which is one-to-one and onto. This result calls into question whether there *could be* a canonical domain in $\mathbb{C}^n$, $n > 1$.

We shall reproduce here Poincaré's original proof. It is remarkable in that it uses the automorphism group of a domain as a biholomorphic invariant.

Let us recall now a few of the elementary ideas from the function theory of several complex variables. Let $\Omega \subseteq \mathbb{C}^n$ be a domain. A function $f : \Omega \rightarrow \mathbb{C}$ is said to be *holomorphic* if it is holomorphic in each variable separately. That is to say, if the values of $z_1, z_2, \dots, z_{j-1}, z_{j+1}, \dots, z_n$ are frozen, then we mandate that $\zeta \mapsto f(z_1, z_2, \dots, z_{j-1}, \zeta, z_{j+1}, \dots, z_n)$ be holomorphic as a function of one complex variable (wherever this function makes sense). It can be shown, thanks to Hartogs (see [KRA3]) that such a function is C^∞ as a function of several variables, indeed it is real analytic as a multi-variable function. Certainly such a function satisfies the Cauchy-Riemann equations.

Now if Ω is a domain then we let the *automorphism group* of Ω , denoted $\text{Aut}(\Omega)$, be the collection of holomorphic maps

$$\Phi(z) = (\varphi_1(z), \dots, \varphi_n(z)) = (\varphi_1(z_1, \dots, z_n), \dots, \varphi_n(z_1, \dots, z_n))$$

from Ω to itself which are both one-to-one and onto. The collection $\text{Aut}(\Omega)$ of such mappings forms a group when equipped with the binary operation of composition of mappings. We use the topology of *uniform convergence on compact sets* (equivalently, the compact-open topology) on the automorphism group. Thus $\text{Aut}(\Omega)$ is a topological group, and in fact (in case Ω is bounded), it can be shown to be a Lie group (see [KOB]).

It is important to note that the automorphism group is a biholomorphic invariant. This means the following. Let

$$\Phi : \Omega_1 \rightarrow \Omega_2$$

be a biholomorphic mapping. Then the induced mapping

$$\text{Aut}(\Omega_1) \ni \varphi \longmapsto \Phi \circ \varphi \circ \Phi^{-1} \in \text{Aut}(\Omega_2)$$

is a one-to-one, surjective group homomorphism (i.e., an *isomorphism*). In particular, if two domains have automorphism groups which are not isomorphic, then the two domains cannot be biholomorphic. This is the strategy that we shall use to show that B and B^n are not biholomorphic. For simplicity, and with no loss of generality, we restrict attention to the case $n = 2$.

We shall leave certain small portions of this argument as exercises for the reader, but all the key ideas will be indicated.

Notice that, for any $\alpha, \beta \in \mathbb{C}$, $|\alpha|, |\beta| < 1$, the mapping

$$\psi_{\alpha, \beta} : (z_1, z_2) \mapsto \left(\frac{z_1 - \alpha}{1 - \overline{\alpha}z_1}, \frac{z_2 - \beta}{1 - \overline{\beta}z_2} \right)$$

is a biholomorphic self-mapping (i.e., an automorphism) of the bidisc D^2 . We assume, seeking a contradiction, that $\Phi : B \rightarrow D^2$ is in fact a biholomorphic mapping of B with D^2 . If $(\alpha, \beta) = \Phi(0, 0)$, then $\psi_{\alpha, \beta} \circ \Phi$ is also a biholomorphic mapping of B to D^2 that takes the origin $(0, 0)$ to the origin $(0, 0)$. For convenience we shall denote this new mapping with the same name Φ .

For any domain Ω with $0 \in \Omega$, we let I_0^Ω denote the *isotropy group* of the origin: this is the subgroup of $\text{Aut}(\Omega)$ consisting of those mappings which fix the origin. Obviously the putative biholomorphic mapping Φ of B to D^2 maps I_0^B in $\text{Aut}(B)$ isomorphically onto $I_0^{D^2}$ in $\text{Aut}(D^2)$. We shall show now that this is impossible.

In fact the connected component of the identity in each of these subgroups will also be mapped isomorphically to each other. So let us consider what those might be. In the bidisc, the maps that fix the origin are (by a simple application of the Schwarz lemma), of two kinds:

- Maps which are just rotations in each variable separately.
- Maps which permute the two variables.

And of course we also must allow compositions of these two types of mappings. But if we are to consider the connected component of the identity in the isotropy group of the origin then we must restrict attention to the first kind. Call the group G_{D^2} . Then we see that

$$G_{D^2} = (\text{maps which are rotations in each variable separately}).$$

Now let us look at the ball B . The maps that fix the origin are of two kinds:

- Unitary rotations.
- Antipodal mappings.

Again, if we wish to consider only the connected component of the identity, then we can only consider mappings of the first kind. Thus

$$G_B = (\text{unitary rotations}).$$

Now the key fact to notice here is that G_{D^2} is *abelian*, while G_B is *not abelian*. Thus these two groups cannot be isomorphic. We conclude then that D^2 and B cannot be biholomorphic. That is Poincaré's theorem.

6 Partial Differential Equations

In the present section we use ideas from Section 2 to derive a formula for the solution of the inhomogeneous Cauchy-Riemann equation. This will be a new idea for most readers. Afterwards we shall supply an interesting application of the ideas.

Recall that the classical Cauchy-Riemann equations for a function $f(z) = u(z) + iv(z)$ are

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

Expressed in terms of the complex derivative, these equations may be written neatly as

$$\frac{\partial}{\partial \bar{z}} f \equiv 0.$$

It turns out to be a matter of considerable interest to solve the equation

$$\frac{\partial}{\partial \bar{z}} f \equiv \alpha$$

for a suitable function α . We shall prove the following result.

Theorem 10 *Let $\alpha \in C_c^1(\mathbb{C})$. The function defined by*

$$f(\zeta) = -\frac{1}{2\pi i} \int \frac{\alpha(\xi)}{\xi - \zeta} d\bar{\xi} \wedge d\xi = -\frac{1}{\pi} \int \frac{\alpha(\xi)}{\xi - \zeta} dA(\xi),$$

where dA is area measure, satisfies

$$\bar{\partial} f(\zeta) = \frac{\partial f}{\partial \bar{\zeta}}(\zeta) d\bar{\zeta} = \alpha(\zeta) d\bar{\zeta}.$$

Proof: Let $D(0, R)$ be a large disc that contains the support of α . Then

$$\begin{aligned}
\frac{\partial f}{\partial \bar{\zeta}}(\zeta) &= -\frac{1}{2\pi i} \frac{\partial}{\partial \bar{\zeta}} \int_{\mathbb{C}} \frac{\alpha(\xi)}{\xi - \zeta} d\bar{\xi} \wedge d\xi \\
&= -\frac{1}{2\pi i} \frac{\partial}{\partial \bar{\zeta}} \int_{\mathbb{C}} \frac{\alpha(\xi + \zeta)}{\xi} d\bar{\xi} \wedge d\xi \\
&= -\frac{1}{2\pi i} \int_{\mathbb{C}} \frac{\frac{\partial \alpha}{\partial \xi}(\xi + \zeta)}{\xi} \delta \bar{\xi} \wedge d\xi \\
&= -\frac{1}{2\pi i} \int_{D(0, R)} \frac{\frac{\partial \alpha}{\partial \xi}(\xi)}{\xi - \zeta} \delta \bar{\xi} \wedge d\xi.
\end{aligned}$$

By Theorem 3, this last equals

$$\alpha(\zeta) - \frac{1}{2\pi i} \int_{\partial D(0, R)} \frac{\alpha(\xi)}{\xi - \zeta} d\xi = \alpha(\zeta).$$

Here we have used the support condition on α . This is the result that we wish to prove. $\square$

A simple limiting argument shows that the theorem remains valid if the function α is only bounded (and not smooth with compact support). We omit the details of the proof. This additional observation will prove useful in the example that follows.

In general the function f that we produce in the last theorem as a solution of the $\bar{\partial}$ equation will *not* be compactly supported—even though α itself *is* compactly supported. See [KRA3] for the details of this assertion. It follows from elementary estimates that the solution f *is* bounded. We now give a simple example of the utility of these partial differential equations ideas.

EXAMPLE 2 Let $\Omega \subseteq \mathbb{C}$ be a domain. Let $P \in \partial\Omega$ and U a small, open neighborhood of P . Suppose that h is a function, holomorphic on $U \cap \Omega$, such that h blows up at P at a prescribed rate. Then there is a holomorphic function $\hat{h}$ on Ω that blows up at P at that same prescribed rate.

To see this, let φ be a C_c^∞ function with support in U that is identically equal to 1 in a neighborhood of P . Define

$$\alpha(z) = \left(\frac{\partial \varphi}{\partial \bar{z}} \right) \cdot h.$$

Then certainly α is a bounded function—for $\partial\varphi/\partial\bar{z}$ is identically 0 in a neighborhood of P . Thus the equation

$$\frac{\partial f}{\partial\bar{z}} = \alpha$$

has a bounded solution. But then the function

$$\hat{h} = \varphi \cdot h - f$$

will satisfy the Cauchy-Riemann equations—since

$$\frac{\partial}{\partial\bar{z}} [\varphi \cdot h - f] = \frac{\partial\varphi}{\partial\bar{z}} \cdot h - \frac{\partial\varphi}{\partial\bar{z}} \cdot h \equiv 0.$$

Moreover, we see that $\hat{h}$ will blow up at P at just the same rate as h itself, since the two functions differ there by the bounded function f . That completes our analysis. $\square$

In this example we have been deliberately nonspecific about what “blows up at a prescribed rate” means. We are endeavoring to avoid the question of whether the singularity is a pole or an essential singularity. In point of fact, this is *not* an isolated singularity in the usual sense of complex variable theory (see [GRK]). Rather, this particular singularity is at the boundary—so the familiar analyses do not apply.

7 Geometry

Ever since Lars Ahlfors’s seminal paper [AHL] on the Schwarz lemma, geometry has played a decisive role in complex function theory. Today it is one of the major tools. In this section we show how Poincaré’s theorem on the biholomorphic inequivalence of the ball and the polydisc (already treated in Section 5 from a different point of view) may be derived using geometric ideas. This is both a validation of the geometric viewpoint and an introduction to some important ideas that grow out of the Riemann mapping theorem.

As we have done in the past, we restrict attention to dimension $n = 2$ for simplicity and convenience. The reader may check that the arguments go through unchanged in all dimensions.

We begin now with some terminology.

Definition 3 If $U_1, U_2 \subseteq \mathbb{C}^2$ and

$$f = (f_1, f_2) : U_1 \longrightarrow U_2$$

is holomorphic then define $\text{Jac}_{\mathbb{C}}f(z)$, $z \in U_1$, (the *Jacobian matrix* of f at P) to be the matrix

$$\begin{pmatrix} \frac{\partial f_1}{\partial z_1}(z) & \frac{\partial f_1}{\partial z_2}(z) \\ \frac{\partial f_2}{\partial z_1}(z) & \frac{\partial f_2}{\partial z_2}(z) \end{pmatrix}$$

Remark 11 Notice that $\text{Jac}_{\mathbb{C}}f(z)$ is distinct from the real Jacobian, $\text{Jac}_{\mathbb{R}}f$, of calculus: the latter would be a 4×4 matrix which arises from treating f as a function from a domain in $\mathbb{R}^4$ to a domain in $\mathbb{R}^4$. $\square$

We continue to use the symbol D to denote the unit disc in $\mathbb{C}$. Let $\Omega \subseteq \mathbb{C}^2$ be a domain and $P \in \Omega$. Define $(D, \Omega)_P$ to be the holomorphic functions $f : \Omega \rightarrow D$ such that $f(P) = 0$. Define $(\Omega, D)_P$ to be the holomorphic functions $f : D \rightarrow \Omega$ such that $f(0) = P$. We are now going to define the Carathéodory and Kobayashi metrics. Since we are working in a two-dimensional space, we can no longer specify a metric as a scalar-valued function on the domain. In fact a metric will measure the length of a vector at a point. Here and throughout, $|\cdot|$ denotes Euclidean length.

Definition 4 If $P \in \Omega$ and $\xi \in \mathbb{C}^2$ define the *Carathéodory length* of ξ at P to be

$$F_C^\Omega(P, \xi) = \sup\{|\text{Jac}_{\mathbb{C}}f(P)\xi| : f \in (D, \Omega)_P\}.$$

Definition 5 If $P \in \Omega$ and $\xi \in \mathbb{C}^2$ define the *Kobayashi length* of ξ at P to be

$$F_K^\Omega(P, \xi) = \inf\{|\xi|/|g'(0)| : g \in (\Omega, D)_P, \\ g'(0) \text{ is a scalar multiple of } \xi\}.$$

The way that we define metrics is motivated by Riemann's philosophy. If $\gamma : [0, 1] \rightarrow \Omega$ is a continuously differentiable curve, we define its Kobayashi length to be

$$\ell_K(\gamma) = \int_0^1 F_K^\Omega(\gamma(t), \gamma'(t)) dt.$$

Notice that we are integrating the lengths, in the metric, of the tangent vectors to the curve. The Carathéodory length of a curve is defined similarly.

One of the basic properties that we shall prove is that holomorphic mappings decrease distance in the Carathéodory and Kobayashi metrics. In several variables we express this assertion as follows.

Proposition 12 *Let U_1 and U_2 be domains and*

$$f : U_1 \longrightarrow U_2$$

be a holomorphic mapping. If $P \in U_1$ and $\xi \in \mathbb{C}^2$ we define

$$f_*(P)\xi = \text{Jac}_{\mathbb{C}}f(P)\xi.$$

Then

$$F_C^{U_1}(P, \xi) \geq F_C^{U_2}(f(P), f_*(P)\xi)$$

and

$$F_K^{U_1}(P, \xi) \geq F_K^{U_2}(f(P), f_*(P)\xi).$$

Proof: We give the proof for the Carathéodory metric. The proof for the Kobayashi metric is similar.

Choose $\varphi \in (D, U_2)_{f(P)}$. Then $\varphi \circ f \in (D, U_1)_P$. Hence

$$\begin{aligned} F_C^{U_1}(P, \xi) &\geq |(\text{Jac}_{\mathbb{C}}(\varphi \circ f)(P))\xi| \\ &= |\text{Jac}_{\mathbb{C}}\varphi(f(P)) \circ (\text{Jac}_{\mathbb{C}}f(P))\xi| \\ &= |\text{Jac}_{\mathbb{C}}\varphi(f(P))(f_*(P)\xi)|. \end{aligned}$$

Taking the supremum over all φ gives

$$F_C^{U_1}(P, \xi) \geq F_C^{U_2}(f(P), f_*(P)\xi). \quad \square$$

Corollary 13 *If f is a biholomorphic map then it preserves both the Carathéodory and the Kobayashi metrics; that is, the inequalities in the proposition become equalities.*

Proof: Obvious. $\square$

Exercise: Check that the proposition implies that f decreases the lengths of curves. That is, if γ is a continuously differentiable curve in U_1 and $f_*\gamma \equiv f \circ \gamma$ is the corresponding curve in U_2 then show that $\ell_C(f_*\gamma) \leq \ell_C(\gamma)$ and $\ell_K(f_*\gamma) \leq \ell_K(\gamma)$. $\square$

We now define two interesting new invariants. These invariants were trivial in one complex variable, but now they provide crucial information.

Definition 6 Let $\Omega \subseteq \mathbb{C}^2$ be a domain and $P \in \Omega$. The *Carathéodory indicatrix* of Ω at P is

$$\mathbf{i}_P^C(\Omega) = \{\xi \in \mathbb{C} : F_C^\Omega(P, \xi) < 1\}.$$

The *Kobayashi indicatrix* of Ω at P is

$$\mathbf{i}_P^K(\Omega) = \{\xi \in \mathbb{C} : F_K^\Omega(P, \xi) < 1\}.$$

In words, the indicatrix is the “unit ball,” in the indicated metric, of vectors at P .

Proposition 14 Let $f : \Omega_1 \rightarrow \Omega_2$ be a biholomorphic mapping of domains in $\mathbb{C}^2$. Say that $f(P) = Q$. Then

$$\text{Jac}_{\mathbb{C}}f(P) : \mathbf{i}_P^C(\Omega_1) \longrightarrow \mathbf{i}_Q^C(\Omega_2)$$

and

$$\text{Jac}_{\mathbb{C}}f(P) : \mathbf{i}_P^K(\Omega_1) \longrightarrow \mathbf{i}_Q^K(\Omega_2)$$

are linear isomorphisms.

Proof: Since f is distance decreasing in the Kobayashi metric, $\text{Jac}_{\mathbb{C}}f(P)$ maps $\mathbf{i}_P^K(\Omega_1)$ into $\mathbf{i}_Q^K(\Omega_2)$. But the same observation applies to

$$\text{Jac}_{\mathbb{C}}(f^{-1})(Q) = (\text{Jac}_{\mathbb{C}}f(P))^{-1};$$

it maps $\mathbf{i}_Q^K(\Omega_2)$ into $\mathbf{i}_P^K(\Omega_1)$. Thus $\text{Jac}_{\mathbb{C}}f(P)$ is a linear isomorphism of $\mathbf{i}_P^K$ to $\mathbf{i}_Q^K$ as claimed.

The proof for $\mathbf{i}_P^C$ is identical. $\square$

Proposition 15 *Let $B = B(0, 1)$ be the unit ball. Then we have*

$$\mathbf{i}_0^K(B) = B.$$

Proof: Let $\varphi \in (B, D)_0$. If η is any Euclidean unit vector in $\mathbb{C}^2$ then consider the function

$$h(\zeta) \equiv \varphi(\zeta) \cdot \eta.$$

Here “ $\cdot$ ” denotes the usual inner product of 2-vectors. We have that h maps the disc to the disc and $h(0) = 0$. By the Schwarz lemma of one variable,

$$|h'(0)| \leq 1.$$

Since this inequality holds for any choice of η , we conclude that

$$|\varphi'(0)| \leq 1.$$

Now if ξ is any vector in $\mathbb{C}^2$ then it follows from the preceding calculation that

$$\begin{aligned} F_K^B(0, \xi) &= \inf\{|\xi|/|\varphi'(0)| : \varphi \in (B, D)_0\} \\ &\geq |\xi|. \end{aligned}$$

On the other hand, the map

$$\varphi_0(\zeta) \equiv \frac{\zeta}{|\xi|}\xi$$

satisfies $\varphi_0 \in (B, D)_0$ and $\varphi'_0(0)$ is a positive multiple of ξ . Therefore

$$F_K^B(0, \xi) \leq |\xi|/|\varphi'_0(0)| = |\xi|.$$

We conclude that

$$F_K^\Omega(0, \xi) = |\xi|,$$

hence that

$$\mathbf{i}_0^K(B) = B.$$

□

Proposition 16 *Let $D^2 = D^2(0, 1)$ be the unit bidisc. Then*

$$\mathbf{i}_0^K(D^2) = D^2.$$

Proof: Define the projections

$$\pi_1(z_1, z_2) = z_1 \quad \text{and} \quad \pi_2(z_1, z_2) = z_2.$$

Let $\eta = (\eta_1, \eta_2) \in \mathbb{C}^2$ be any vector. By Proposition 3, we have that

$$F_K^{D^2}(0, \eta) \geq F_K^{\pi_1(D^2)}(\pi_1(0), (\pi_1)_*\eta) = F_K^D(0, \eta_1).$$

But the Schwarz lemma of one variable tells us easily that the last quantity is just $|\eta_1|$. A similar argument shows that

$$F_K^{D^2}(0, \eta) \geq |\eta_2|.$$

We conclude from these two inequalities that

$$F_K^{D^2}(0, \eta) \geq \max\{|\eta_1|, |\eta_2|\}.$$

Therefore

$$\mathbf{i}_0^K(D^2) \subseteq D^2.$$

For the reverse inclusion, fix η as above and consider the function

$$\varphi(\zeta) = \left(\frac{\zeta \eta_1}{\max\{|\eta_1|, |\eta_2|\}}, \frac{\zeta \eta_2}{\max\{|\eta_1|, |\eta_2|\}} \right).$$

Then it is obvious that $\varphi \in (\Omega, D)_0$ and that $\varphi'(0)$ is a positive multiple of η . Therefore

$$F_K^\Omega(0, \eta) \leq \frac{|\eta|}{|\varphi'(0)|} = \max\{|\eta_1|, |\eta_2|\}.$$

The opposite inclusion now follows. □

Exercise: Verify that $\mathbf{i}_0^C(B) = B$ and $\mathbf{i}_0^C(D^2) = D^2$. □

Theorem 17 (Poincaré) *There is no biholomorphic mapping of the bidisc D^2 to the ball B .*

Proof: Suppose that

$$\Phi : D^2 \longrightarrow B$$

is biholomorphic. Let $\Phi^{-1}(0) = \alpha \in D^2$. There is an element $\varphi \in \text{Aut}(D^2(0, 1))$ such that $\varphi(0) = \alpha$. Consider $g \equiv \Phi \circ \varphi$. Then

$$g : D^2 \longrightarrow B$$

is biholomorphic and $g(0) = 0$. We will show that g cannot exist.

By Proposition 11, $\text{Jac}_{\mathbb{C}}g(0)$ is a linear isomorphism of $\mathbf{i}_0^K(D^2)$ to $\mathbf{i}_0^K(B)$. But Propositions 12 and 13 identify these as D^2 and B respectively. So we have that

$$\text{Jac}_{\mathbb{C}}g(0) : D^2 \longrightarrow B$$

is a linear isomorphism. However this is impossible. For the segment $\mathbf{i} = \{(t + i0, 1) : 0 \leq t \leq 1\}$ lies in ∂D^2 . The linear isomorphism would map $\mathbf{i}$ to a nontrivial segment in ∂B . But B is strictly convex (all boundary points are extreme) so its boundary contains no segments. This is the required contradiction. $\square$

Remark 18 It is important to appreciate the logic in this proof. The hypothesized map Φ (and therefore g as well) is *not* assumed to extend in any way to ∂D^2 . Indeed, given the very different natures of ∂D^2 and ∂B , we would expect Φ to be highly pathological at the boundary. Our geometric machinery allows us to pass to the linear map $\text{Jac}_{\mathbb{C}}g(0)$, which is defined on all of space. Thus we are able to analyze the boundaries of the domains and arrive at a contradiction. $\square$

8 Concluding Remarks

We have given several examples to show how a problem that began as a complex analysis problem could ultimately be resolved by techniques from another part of mathematics. This is not to say that one could not have brought function-theoretic techniques to bear on the original problems. But it is always fruitful to bring in new ideas and new perspectives.

In fact it is safe to say that the application of techniques from other disciplines has helped to spawn entirely new directions in mathematics. Today the subjects of Kähler geometry, the $\bar{\partial}$ -Neumann problem, and automorphism

groups—to give just a few examples—are branches of complex analysis that have a life of their own.

Moreover, those new ideas will often bear additional benefits. They will give new insights, offer different points of view, or (in the best of all possible circumstances) suggest new problems. Certainly that is the way that matters have played out in practice in modern complex function theory.

Other parts of mathematics have seen similar benefits from bringing in ideas from the outside. In recent years we have seen applications of techniques from partial differential equations in algebraic geometry, applications of Lie group theory in several complex variables, and applications of abstract logic in real analysis. The number of examples keeps growing, and mathematics is correspondingly enriched. We hope that this article has served as a modest introduction to this exciting set of ideas.

REFERENCES

- [AHL] L. Ahlfors, An extension of Schwarz's lemma *Trans. Amer. Math. Soc.* 43 (1938), 359–364.
- [BLK] B. Blank and S. G. Krantz, *Calculus, Multivariable*, Key College Press, Emeryville, CA, 2006.
- [GIT] Gilbarg and N. Trudinger, *Elliptic Partial Differential Equations of Second Order*, 2nd ed., Springer-Verlag, Berlin, 2001.
- [GRK] R. E. Greene and S. G. Krantz, *Function Theory of One Complex Variable*, 3rd ed., American Mathematical Society, Providence, RI, 2006.
- [HOR] L. Hörmander, *Linear Partial Differential Equations*, Springer-Verlag, Berlin, 1969.
- [KOB] S. Kobayashi, *Hyperbolic Manifolds and Holomorphic Mappings*, Marcel Dekker, New York, 1970.
- [KRA1] S. G. Krantz, The Carathéodory and Kobayashi metrics and applications in complex analysis, *American Mathematical Monthly*, to appear.
- [KRA2] S. G. Krantz, *Partial Differential Equations and Complex Analysis*, CRC Press, Boca Raton, FL, 1992.
- [KRA3] S. G. Krantz, *Function Theory of Several Complex Variables*, 2nd ed., American Mathematical Society, Providence, RI, 2001.
- [KRA4] S. G. Krantz, *Geometric Function Theory: Exploration in Complex Analysis*, Birkhäuser, Boston, 2006.
- [KRA5] S. G. Krantz, *A Panorama of Harmonic Analysis*, Mathematical Association of America, Washington, D.C., 1999.
- [SIC] J. Siciak, On series of homogeneous polynomials and their partial sums, *Annales Polonici Math.* 51(1990), 289–302.
- [STE] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, NJ, 1970.

STEVEN G. KRANTZ received his B.A. degree from the University of California at Santa Cruz in 1971. He earned the Ph.D. from Princeton University in 1974. He has taught at UCLA, Princeton University, Penn State, and Washington University in St. Louis. Krantz is the holder of the UCLA Alumni Foundation Distinguished Teaching Award, the Chauvenet Prize, and the Beckenbach Book Prize. He is the author of 150 papers and 50 books. His research interests include complex analysis, real analysis, harmonic analysis, and partial differential equations. Krantz is currently the Deputy Director of the American Institute of Mathematics.

*American Institute of Mathematics, 360 Portage Avenue,
Palo Alto, CA 94306*

sk@math.wustl.edu